arXiv:1106.3476v1 [math.CV] 17 Jun 2011

THE RATE OF INCREASE OF MEAN VALUES OF FUNCTIONS IN WEIGHTED HARDY SPACES

CHENGJI XIONG & JUNMING LIU

Abstract. Let $0 < p < \infty$ and $0 \leq q < \infty$. For each f in the weighted Hardy space $H_{p,q}$, we show that $d\|f_r\|_{p,q}^p/dr$ grows at most like $o(1/1-r)$ as $r \to 1$.

1. Introduction

Let $\mathbb{D}$ be the unit disk in the complex plane and $0 < p < \infty$. The Hardy space $H^p(\mathbb{D})$ is the family of all analytic functions f in $\mathbb{D}$ satisfying

$$\|f\|_p = \lim_{r\to 1} \|f_r\|_p < \infty,$$

where

$$\|f_r\|_p = \Big(\frac{1}{2\pi}\int_0^{2\pi} |f(re^{i\theta})|^p d\theta\Big)^{1/p}.$$

If $0 \leq q < \infty$, define

$$\|f_r\|_{p,q} = \Big(\frac{1}{2\pi}\int_0^{2\pi} |f(re^{i\theta})|^p (1-r^2)^q d\theta\Big)^{1/p}.$$

The weighted Hardy space $H_{p,q}(\mathbb{D})$ is the family of all analytic functions f in $\mathbb{D}$ satisfying

$$\|f\|_{p,q} = \sup_{0<r<1} \|f_r\|_{p,q} < \infty.$$

It is trivial that $H_{p,0}$ is just the classical Hardy space H^p and $H_{p,q}$ is a Banach space for $p \geq 1$. There are many books about H^p and $H_{p,q}$ such as Duren's book [3] and Zhu's book [7]. It is well known that $\|f_r\|_p$ is a nondecreasing function of r. Furthermore, Hardy [4] showed that $\log \|f_r\|_p$ is a convex function of $\log r$. Other properties of mean values of analytic functions can be found in [1, 2, 6].

Recently, Mashreghi [5] showed that $d\|f_r\|_p/dr$ grows at most like $o(1/1-r)$ as $r \to 1$. In this paper, we generalized this result to weighted Hardy spaces.

2. Lemmas

Suppose that $f \not\equiv 0$ is analytic in $\mathbb{D}$. Let $W(z) = |f(z)|^p(1-|z|^2)^q$ for $0 < p < \infty$, $0 \leq q < \infty$, and ∇ be the gradient operator.

Lemma 1. *Let $0 < r \leq 1$ and $D_r = \{z : |z| < r\}$. For $z_0 \in D_r$ and small $\varepsilon > 0$, let $D(z_0,\varepsilon) = \{z : |z - z_0| \leq \varepsilon\} \subset D_r$ and*

$$I_\varepsilon = \int_{\partial D(z_0,\varepsilon)} \Big(\log(r/|z|)\frac{\partial W}{\partial n} - W\frac{\partial \log(r/|z|)}{\partial n}\Big) d\ell.$$

If $z_0 \neq 0$ is a zero of f, then $\lim_{\varepsilon\to 0} I_\varepsilon = 0$. If $z_0 = 0$ then $\lim_{\varepsilon\to 0} I_\varepsilon = 2\pi|f(0)|^p$.

2000 *Mathematics Subject Classification.* 30H10.

Key words and phrases. Hardy space, weighted Hardy space, mean value.

Proof. Since $W(z)=|f(z)|^p(1-|z|^2)^q$, direct computation gives

$$\nabla W(z)=(1-|z|^2)^q\nabla|f(z)|^p+|f(z)|^p\nabla(1-|z|^2)^q$$

and

$$|\nabla|f(z)|^p|\le p|f(z)|^{p-1}|f'(z)|,\quad |\nabla(1-|z|^2)^q|=2q|z|(1-|z|^2)^{q-1}.$$

So

$$\begin{aligned}\Big|\frac{\partial W(z)}{\partial n}\Big|&\le\big|\nabla W(z)\big|\le(1-|z|^2)^q|\nabla|f(z)|^p|+|f(z)|^p|\nabla(1-|z|^2)^q|\\&\le p|f(z)|^{p-1}|f'(z)|(1-|z|^2)^q+2q|z||f(z)|^p(1-|z|^2)^{q-1}\end{aligned}$$

and

$$\Big|\frac{\partial\log(r/|z|)}{\partial n}\Big|\le|\nabla\log(r/|z|)|=\frac{1}{|z|}.$$

Write

$$\begin{aligned}I_\varepsilon&=\int_{\partial D(z_0,\varepsilon)}\Big(\log(r/|z|)\frac{\partial W}{\partial n}-W\frac{\partial\log(r/|z|)}{\partial n}\Big)d\ell\\&=\int_0^{2\pi}\log(r/|z_0+\varepsilon e^{i\theta}|)\frac{\partial W(z_0+\varepsilon e^{i\theta})}{\partial n}\varepsilon d\theta-\int_{\partial D(z_0,\varepsilon)}W(z)\frac{\partial\log(r/|z|)}{\partial n}d\ell\\&=I_1-I_2.\end{aligned}$$

For convenience, let $C>0$ be a constant independent of ε, whose value may change from line to line.

If $z_0\neq0$ is a zero of order $k\ge1$, then $|I_1|\le C\varepsilon^{kp}$ and $|I_2|\le C\varepsilon^{k+1}$. Thus $\lim_{\varepsilon\to0}I_\varepsilon(z_0)=0$.

If $z_0=0$ and $f(z_0)\neq0$, then $|I_1|\le\varepsilon\log(r/\varepsilon)$ and

$$-I_2=\int_0^{2\pi}W(\varepsilon e^{i\theta})d\theta=(1-\varepsilon^2)^q\int_0^{2\pi}|f(\varepsilon e^{i\theta})|^pd\theta.$$

Hence $\lim_{\varepsilon\to0}I_\varepsilon=2\pi|f(0)|^p$.

At last, if $z_0=0$ is a zero of f of order $k\ge1$, then $|I_1|\le C\varepsilon^{kp}\log(r/\varepsilon)$ and $|I_2|\le C\varepsilon^{kp}$. We have $\lim_{\varepsilon\to0}I_\varepsilon=2\pi|f(0)|^p$ also. □

Lemma 2. *Let $f\in H_{p,\ q}(\mathbb{D})$, $0<p<\infty$, $0\le q<\infty$, and $f\not\equiv0$. Then*

$$\lim_{r\to1}\int_0^{2\pi}|f(re^{i\theta})|^p(1-r^2)^qd\theta-2\pi|f(0)|^p=\int_{|z|<1}\log(1/|z|)G(z)dxdy\ ,$$

where

$$G(z)=(1-|z|^2)^q\nabla^2|f|^p+2\nabla|f|^p\cdot\nabla(1-|z|^2)^q+|f|^p\nabla^2(1-|z|^2)^q.$$

Proof. Since $f\not\equiv0$ is analytic in $\mathbb{D}$, for any $0<R<1$, we can choose r, $R<r<1$ such that f has finite many zeros in $D_r=\{|z|<r\}$ and no zeros on the circle ∂D_r. Let $\{z_1,z_2,\cdots,z_n\}$ be the set consisting of 0 and all zeros of f in D_r. Take $\varepsilon>0$ so small that $\{z:|z-z_k|\le\varepsilon\}\subset D_r$ for $1\le k\le n$, and all these disks are disjoint. Denote

$$\Omega=D_r\backslash\bigcup_{k=1}^n\{z:|z-z_k|\le\varepsilon\}.$$

The function $W(z) = |f(z)|^p(1-|z|^2)^q$ is infinitely differentiable in a neighborhood of $\overline{\Omega}$. Then by Green's theorem, we have

$$\int_\Omega \log(r/|z|)\nabla^2 W(z)dxdy = \int_{\partial\Omega}\Big(\log(r/|z|)\frac{\partial W}{\partial n} - W\frac{\partial \log(r/|z|)}{\partial n}\Big)d\ell. \tag{2.1}$$

Direct computation gives

$$\nabla^2 W(z) = (1-|z|^2)^q\nabla^2|f|^p + 2\nabla|f|^p\cdot\nabla(1-|z|^2)^q + |f|^p\nabla^2(1-|z|^2)^q = G(z).$$

The direction of every small circle in $\partial\Omega$ is clockwise. For $\varepsilon \to 0$, by Lemma 1, formula (2.1) becomes

$$\int_{\partial D_r}\Big(\log(r/|z|)\frac{\partial W}{\partial n} - W\frac{\partial \log(r/|z|)}{\partial n}\Big)d\ell - 2\pi|f(0)|^p = \int_{D_r}\log(r/|z|)G(z)dxdy.$$

The left-side of above formula is equal to

$$\int_0^{2\pi} W(re^{i\theta})d\theta - 2\pi|f(0)|^p.$$

Now we obtain the lemma as $r \to 1$. □

3. The rate of increase of $\|f_r\|_{p,q}^p$

Theorem 1. *Let* $f \in H_{p,\ q}(\mathbb{D})$ *and* $f \not\equiv 0$, *then*

$$\frac{d\|f_r\|_{p,q}^p}{dr} = o(1/1-r) \text{ as } r \to 1.$$

Proof. As in the proof of Lemma 2, we choose suitable r and ε. Replacing $\log r/|z|$ by $\log 1/|z|$, by Green's theorem, we have,

$$\int_\Omega \log(1/|z|)\nabla^2 W(z)dxdy = \int_{\partial\Omega}\Big(\log(1/|z|)\frac{\partial W}{\partial n} - W\frac{\partial \log(1/|z|)}{\partial n}\Big)d\ell. \tag{3.1}$$

For $\varepsilon \to 0$, using Lemma 1 for $r = 1$, formula (3.1) turns into

$$\begin{aligned}\int_{D_r}\log(1/|z|)G(z)dxdy &= \int_{\partial D_r}\Big(\log(1/|z|)\frac{\partial W}{\partial n} - W\frac{\partial \log(1/|z|)}{\partial n}\Big)d\ell - 2\pi|f(0)|^p\\ &= \int_0^{2\pi}\Big(\log\frac{1}{r}\frac{\partial W}{\partial r}(re^{i\theta}) + \frac{W(re^{i\theta})}{r}\Big)rd\theta - 2\pi|f(0)|^p\\ &= \int_0^{2\pi} W(re^{i\theta})d\theta - r\log r\int_0^{2\pi}\frac{\partial W}{\partial r}(re^{i\theta})d\theta - 2\pi|f(0)|^p\\ &= \int_0^{2\pi}|f(re^{i\theta})|^p(1-r^2)^q d\theta - 2\pi r\log r\frac{d\|f_r\|_{p,q}^p}{dr} - 2\pi|f(0)|^p.\end{aligned}$$

Now let $r \to 1$. By Lemma 2, we have

$$\lim_{r\to 1}\log r\frac{d\|f_r\|_{p,q}^p}{dr} = 0.$$

Hence,

$$\lim_{r\to 1}\frac{d\|f_r\|_{p,q}^p}{dr} = o(1/\log r) = o(1/1-r).$$

□

By Theorem 1, if $\lim_{r\to 1}\|f_r\|_{p,q}\neq 0$, or for $q=0$, $\|f_r\|_{p,q}$ is increasing with r, then we can get $\lim_{r\to 1} d\|f_r\|_{p,q}/dr = o(1/1-r)$.

In the proof of Theorem 1, using 1 instead of $\log 1/|z|$, similar arguments can deduce

$$2\pi r\frac{d\|f_r\|_{p,q}^p}{dr}=\int_{|z|<r}G(z)dxdy.$$

Corollary 1. *If $f\in H_{p,q}$ with $0<p<\infty,\ 0\le q<\infty$, then we have*

$$2\lim_{r\to 1}\int_0^{2\pi}|f(re^{i\theta})|^p(1-r^2)^q d\theta=\int_{\mathbb{D}}(1-|z|^2)\nabla^2\Big(|f(z)|^p(1-|z|^2)^q\Big)dxdy$$

$$+\,4\int_{\mathbb{D}}|f(z)|^p(1-|z|^2)^q dxdy. \tag{3.2}$$

Proof. Denote

$$J_\varepsilon=\int_{\partial D(z_0,\varepsilon)}\Big((1-|z|^2)\frac{\partial W}{\partial n}-W\frac{\partial(1-|z|^2)}{\partial n}\Big)d\ell,$$

where z_0 is a zero of f. Similar arguments as in the proof of Lemma 1, we have $\lim_{\varepsilon\to 0}J_\varepsilon=0$. Choosing suitable r and ε as in the proof of Theorem 1, by Green's theorem, we have

$$\int_\Omega(1-|z|^2)\nabla^2W(z)-\nabla^2(1-|z|^2)\cdot W(z)dxdy=\int_{\partial\Omega}\Big((1-|z|^2)\frac{\partial W}{\partial n}-W\frac{\partial(1-|z|^2)}{\partial n}\Big)d\ell$$

As $\varepsilon\to 0$, the above formula turns into

$$\int_{D_r}(1-|z|^2)\nabla^2W(z)+4W(z)dxdy=\int_{\partial D_r}\Big((1-|z|^2)\frac{\partial W}{\partial n}-W\frac{\partial(1-|z|^2)}{\partial n}\Big)d\ell$$

$$=\int_0^{2\pi}\Big((1-r^2)\frac{\partial W}{\partial r}(re^{i\theta})+2rW(re^{i\theta})\Big)rd\theta. \tag{3.3}$$

By Theorem 1, we have

$$\lim_{r\to 1}\int_0^{2\pi}r(1-r^2)\frac{\partial W}{\partial r}(re^{i\theta})d\theta=2\lim_{r\to 1}\left[(1-r)\frac{\partial}{\partial r}\int_0^{2\pi}W(re^{i\theta})d\theta\right]=0\ .$$

The proof is completed by letting $r\to 1$ in formula (3.3). □

Note that $\nabla^2|f|^p=p^2|f|^{p-2}|f'|^2$. Taking $q=0$ in formula (3.2), we obtain

$$\int_0^{2\pi}|f(e^{i\theta})|^p d\theta=\frac{p^2}{2}\int_{\mathbb{D}}(1-|z|^2)|f(z)|^{p-2}|f'(z)|^2dxdy+2\int_{\mathbb{D}}|f(z)|^p dxdy,$$

which is reminiscent of Hardy-Stein identity

$$\frac{1}{2\pi}\int_0^{2\pi}|f(e^{i\theta})|^p d\theta=|f(0)|^p+\frac{p^2}{2\pi}\int_{\mathbb{D}}\log\frac{1}{|z|}|f(z)|^{p-2}|f'(z)|^2dxdy.$$

References

[1] E. F. Beckenbach, *A property of mean values of an analytic function*, Rend. Circ. Mat. Palermo, 1 (1952) no.2, 157-163.

[2] E. F. Beckenbach; W. Gustin and H. Shniad, *On the mean modulus of an analytic function*, Bull. Amer. Math. Soc, 55, (1949) 184-190.

[3] P. L. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics, Vol. 38 Academic Press, New York-London 1970.

[4] G. H. Hardy, *The mean value of the modulus of an analytic function*, Proc. London Math. Soc. 14 (1915), no.1, 269-277.

[5] J. Mashreghi, *The Rate of Increase of Mean Values of Functions in Hardy Spaces*, J. Aust. Math. Soc, 86 (2009), 199-204.

[6] H. Shniad, *Convexity properties of integral means of analytic functions*, Pacific J. Math, 3, (1953), 657-666.

[7] K. Zhu, *Operator theory in function spaces*, Second edition. Mathematical Surveys and Monographs, 138. American Mathematical Society, Providence, RI, 2007.

Department of Mathematics, Shantou University, Shantou, Guangdong, 515063, P. R. China

E-mail address: `chengji_xiong@yahoo.com, 08jmliu@stu.edu.cn`